

A note on De Moivre's quintic equation

M. L. Glasser

Department of Physics, Clarkson University
Potsdam, NY 13699-5820 (USA)

The quintic equation with real coefficients

$$x^5 + 5ax^3 + 5a^2x + b = 0$$

is solved in radicals and the result is used to sum a hypergeometric series for several arguments..

We start with the general quintic equation (over the real numbers)

$$\sum_{j=0}^5 a_j x^j = 0 \quad (1)$$

with $a_5 = 1$. Using Vietá's substitution, $x = u - a/u$ with $a \neq 0$ real, expanding and multiplying by u^5 , the resulting 10-th degree equation is solvable (in radicals) only for $a_1 = aa_3 = 5a^2$, $a_2 = a_4 = 0$, in which case

$$u^{10} + a_0 u^5 - a^5 = 0. \quad (2)$$

By solving this quadratic equation (in u^5) and extracting the fifth roots, after eliminating duplicates, we see that the roots of the De Moivre equation (writing b in place of a_0)

$$x^5 + 5ax^3 + 5a^2x + b = 0 \quad (3)$$

are

$$\begin{aligned} x_1 &= 2^{-1/5}[(\sqrt{\gamma} - b)^{1/5} - (\sqrt{\gamma} + b)^{1/5}] \\ x_2 &= -2^{-1/5}[e^{\pi i/5}(\sqrt{\gamma} - b)^{1/5} + e^{4\pi i/5}(\sqrt{\gamma} + b)^{1/5}] = x_4^* \\ x_3 &= 2^{-1/5}[e^{2\pi i/5}(\sqrt{\gamma} - b)^{1/5} + e^{3\pi i/5}(\sqrt{\gamma} + b)^{1/5}] = x_5^* \\ \gamma &= 4a^5 + b^2. \end{aligned} \quad (4)$$

(The asterisk denotes the complex conjugate.)

By means of the Tschirnhausen transformation [1] $y = x^2 + \alpha x + \beta$ any quintic can be reduced to "principal" form $y^5 + Ay^2 + By + C = 0$. This can be carried out efficiently using Mathematica by means of the command

Resultant $[x^5 + 5ax^3 + 5a^2x + b, y - (x^2 + \alpha x + \beta), x]$ which returns $y^5 + (10a - 5\beta)y^4 + (35a^2 + 5a\alpha^2 - 40a\beta + 10\beta^2)y^3 + \dots$ from which we easily determine $\alpha = \sqrt{a}$ and $\beta = 2a$. Thus, the principal form for the De Moivre quintic is

$$y^5 + (5\sqrt{a}b - 10a^3)y^2 + 15a^4y - (9a^{5/2}b + 22a^5 + b^2) = 0. \quad (5)$$

Jerrard[1] found that by means of a quartic Tschirnhausen transformation a principal quintic can be further reduced to the trinomial Bring form

$$z^5 - z + t = 0. \quad (6)$$

The details can be mechanized to some extent[2], but are still quite cumbersome. To present the result it is convenient to introduce the further abbreviations: $\alpha = 8a^{5/2} - b$, $\beta = 2a^{5/2} - b$, $\delta = 176a^5 + 36a^{5/2}b - b^2$. Also $d_0 = 675(\beta/\alpha)(2a^{5/2} + b)a^2b$, $d_1 = \Delta/\alpha$, $d_2 = 25(\beta/\alpha)(8a^5 - 3a^{5/2}b - b^2)$, $d_3 = 75(\beta/\alpha)a^{3/2}(16a^{5/2} + 3b)$ and $d_4 = 75\beta a^{1/2}$, where

$$\Delta = 25[\frac{\delta\gamma^{1/2}\alpha}{3^{2/3}a^{1/2}}(\beta\sqrt{3a^{3/2}(11a^{5/2} - b)} - 9a^2\gamma^{1/2})^{1/3} +$$

$$\frac{1}{3^{1/3}\alpha\gamma^{1/2}} \frac{16a^{15/2} + 4a^5b + 4a^{5/2}b^2 + b^3}{(\beta\sqrt{3a^{3/2}(11a^{5/2} - b)} - 9a^2\gamma^{1/2})^{1/3}} - (800a^{17/2} - 318a^6b + 227a^{7/2}b^2 - 12ab^3)]. \quad (7)$$

The quantity Δ has six possible values, due to the choice of branches in the square and cube roots, so it takes a bit of experimentation to select the appropriate one.

Next, we require the coefficients

$$c_0 = \frac{5^6\delta^3\gamma^2}{\alpha^5} [5625\beta^2 f_1(a, b) + \frac{25}{a}\beta\Delta f_2(a, b) + 3a^{1/2}\Delta^2(188a^5 + 86a^{5/2}b + 9b^2)] \quad (8a)$$

$$c_1 = \frac{5^5\delta^2\gamma a^{1/2}}{\alpha^4} [5625\beta^2 g_1(a, b) + \frac{25}{a}\beta\Delta g_2(a, b) + 9a^3\Delta^2(4a^{5/2} + b)], \quad (8b)$$

where

$$\begin{aligned} f_1(a, b) &= 2622464a^{35/2} - 1339776a^{15}b + 103828a^{25/2}b^2 + 218210a^{10}b^3 - \\ &\quad 17365a^{15/2}b^4 - 2858a^5b^5 + 297a^{5/2}b^6 - b^7 \\ f_2(a, b) &= 209024a^{25/2} - 30616a^{10}b - 34508a^{15/2}b^2 - \\ &\quad 98a^5b^3 + 530a^{5/2}b^4 - b^5, \\ g_1(a, b) &= 133120a^{35/2} - 83520a^{15}b + 26804a^{25/2}b^2 + 1789a^{10}b^2 - \\ &\quad 1082a^{15/2}b^4 + 386a^5b^5 - 48a^{5/2}b^6 + b^7, \\ g_2(a, b) &= 11200a^{25/2} - 3656a^{10}b - 754a^{15/2}b^2 + 62a^5b^3 - 38a^{5/2}b^4 + b^5. \end{aligned} \quad (9)$$

The result is that the roots of the Bring-Jerrard quintic

$$z^5 - z + t = 0 \quad (10)$$

with

$$t = -e^{-i\pi/4}c_0c_1^{-5/4} \quad (11)$$

are

$$z_j = e^{-i\pi/4}c_1^{-1/4} \sum_{k=0}^4 d_k(x_j^2 + \sqrt{a}x_j + 2a)^k. \quad (12).$$

Now it has been shown by James Cockle[3] (the first Chief Magistrate of Queensland) and others[4] that one of the roots of (10) is

$$z_0 = t {}_4F_3(1/5, 2/5, 3/5, 4/5; 1/2, 3/4, 5/4; \frac{3125}{256}t^4). \quad (13)$$

Therefore, it has been shown that

$${}_4F_3(1/5, 2/5, 3/5, 4/5; 1/2, 3/4, 5/4; -\frac{3125 c_0^4}{256 c_1^5}) = -\frac{c_1}{c_0} \sum_{k=0}^4 d_k (x_0^2 + \sqrt{a}x_0 + 2a)^k \quad (14)$$

where x_0 is one of the five values in (4). (Which one it is can be determined numerically).

As a very simple example, let us take the case $b = 2a^{5/2}$. Then the principal quintic (5) is already in Bring form, which is easily scaled by $y = e^{i\pi/4} 15^{1/4} az$ into (6) with $t = 44e^{-i\pi/4} 15^{-5/4}$. Therefore, from (13) we find

$${}_4F_3(1/5, 2/5, 3/5, 4/5; 1/2, 3/4, 5/4; -\frac{14641}{243}) = \frac{15}{44} [(\sqrt{2} + 1)^{2/5} + (\sqrt{2} - 1)^{2/5} - (\sqrt{2} + 1)^{1/5} + (\sqrt{2} - 1)^{1/5}]. \quad (15)$$

At the other extreme, the algorithm described here [5] produces formulas such as

$${}_4F_3(1/5, 2/5, 3/5, 4/5; 1/2, 3/4, 5/4; z) = Z$$

where

$$z = ((13 + \sqrt{182})^{2/3} ((78913^{2/3} + 24713^{1/6} \sqrt{14})(13 + \sqrt{182})^{1/3} + 2275(13 + \sqrt{182})^{2/3} - 13^{1/3}(6799 + 542\sqrt{182}))^4) / (1521((-713^{2/3} + 313^{1/6} \sqrt{14})(13 + \sqrt{182})^{1/3} - 161(13 + \sqrt{182})^{2/3} + 13^{1/3}(133 + 10\sqrt{182}))^5)$$

and

$$\begin{aligned} Z = & (5((-713^{2/3} + 313^{1/6} \sqrt{14})(13 + \sqrt{182})^{1/3} - 161(13 + \sqrt{182})^{2/3} + 13^{1/3}(133 + 10\sqrt{182}))(-3(-5 + \sqrt{26})^{4/5}(13 + \sqrt{182})^{1/3}(1103 + 3465(5 + \sqrt{26})^{1/5} + 5355(5 + \sqrt{26})^{2/5} + 3780(5 + \sqrt{26})^{3/5}) + 3(-5 + \sqrt{26})^{3/5}(13 + \sqrt{182})^{1/3}(8872 + 1485\sqrt{26} + 4412(5 + \sqrt{26})^{1/5} + 6930(5 + \sqrt{26})^{2/5} + 3780(5 + \sqrt{26})^{4/5}) + (-5 + \sqrt{26})^{1/5}(-14213^{2/3} + 14213^{1/3}(13 + \sqrt{182})^{2/3} + (13 + \sqrt{182})^{1/3}(34114 + 5355\sqrt{26} + 13236(5 + \sqrt{26})^{3/5} + 10395(5 + \sqrt{26})^{4/5} - \end{aligned}$$

$$\begin{aligned}
& 9(5 + \sqrt{26})^{2/5}(-2797 + 180\sqrt{26})) - (-5 + \sqrt{26})^{2/5}(-14213^{2/3} + \\
& 14213^{1/3}(13 + \sqrt{182})^{2/3} + (13 + \sqrt{182})^{1/3}(33742 + 6480\sqrt{26} + \\
& 20790(5 + \sqrt{26})^{3/5} + 16065(5 + \sqrt{26})^{4/5} + \\
& 9(5 + \sqrt{26})^{1/5}(2797 + 180\sqrt{26}))) + (5 + \sqrt{26})^{1/5}(14213^{2/3} - 3309(5 + \sqrt{26})^{3/5}(13 + \sqrt{182})^{1/3} + \\
& 3(5 + \sqrt{26})^{2/5}(-8872 + 1485\sqrt{26})(13 + \sqrt{182})^{1/3} + \\
& (-34114 + 5355\sqrt{26})(13 + \sqrt{182})^{1/3} - 14213^{1/3}(13 + \sqrt{182})^{2/3} + \\
& 2(5 + \sqrt{26})^{1/5}(7113^{2/3} + (-16871 + 3240\sqrt{26})(13 + \sqrt{182})^{1/3} - 7113^{1/3}(13 + \\
& \sqrt{182})^{2/3}))) / (568(13 + \sqrt{182})^{1/3}((78913^{2/3} + \\
& 24713^{1/6}\sqrt{14})(13 + \sqrt{182})^{1/3} + 2275(13 + \sqrt{182})^{2/3} - 13^{1/3}(6799 + 542\sqrt{182})))
\end{aligned}$$

Acknowledgement I am grateful to Prof. Victor Adamchik for help with the necessary Mathematica routines.

References

- [1] V.S. Adamchik and D.J. Jeffrey, ACM SIGSAM Bulletin**37**, 90 (2003).
- [2] Victor Adamchik, Private communication.
- [3] J. Cockle, Phil. Mag.**20**, 145 (1860).
- [4] M.L. Glasser, J. Comp. Appl. Math., **118**, 169 (2000).
- [5] The author has put together a Mathematica notebook incorporating this procedure which will be provided on request to laryg@clarkson.edu.